

FORAGING BEHAVIOUR OF HONEYBEE ON PARENTAL LINES OF HYBRID CAULIFLOWER PUSA HYBRID- 2

P. SELVAKUMAR, S.N. SINHA, V.K. PANDITA, R.M. SRIVASTAVA

Indian Agricultural Research Institute, Regional Station, Karnal, Haryana, INDIA

Abstract

Studies on insect pollination on parental lines of hybrid cauliflower Pusa hybrid 2 revealed that honeybees were predominant pollinators and constituted 85.23%. Pollen gathering honeybees outnumbered nectar collectors. Among honeybees, *Apis dorsata* F., *Apis mellifera* L., *Apis cerana indica* F. and *Apis florea* F. constituted 28.23%, 26.32%, 24.20% and 21.23%, respectively. The number of pollen gatherers reached to its peak at 14.00 hours, while nectar collectors remained constant throughout the day. Foraging behaviour of honeybees on parental lines with regard to their attractiveness showed no significant difference. Frequency of honeybee movement between parental lines viz. male to male, male to female, female to female and female to male also showed no significant difference among them.

Introduction

In cauliflower, F₁ hybrids have been found advantageous for earliness, high yield, bigger curd size, better curd quality, uniform maturity and disease resistance. For large-scale hybrid seed production, self-incompatibility mechanism is being employed. Hybrid seed is obtained by growing together two different inbreds as parental lines and relying mainly upon insect pollinators to cross-pollinate the flowers. Population of natural pollinators differ from location to location depending upon the availability of suitable natural habitat such as forestry and weed flora, which serve as alternate source of nectar and pollen. Investigation on insect pollinators visiting seed production area from natural abode helps to estimate its population. In case of lack of natural pollinators, it is possible to utilise honeybees as they can be reared and their population can be manipulated according to the need of hour. The effectiveness of the honeybees as pollinators for cauliflower seed production has been studied by different workers in open pollinated varieties (RAULA, 1972, SHARMA et al., 1974, ADLAKHA and DHALIWAL, 1979, KAKAR, 1981). Some reports on hybrid seed production in Brussels sprout (FAULKNER, 1974 and 1976) revealed that honeybees were not effective as they were highly selective in their visits to parental lines. Selfed and sibbed seeds in hybrid seed lot were mainly due to selective movement of honeybees. Selfed and sibbed seeds produce plants, which lack vigour and give reduced yields. Hence present study was undertaken to study the foraging behaviour of honeybees on parental lines of hybrid cauliflower Pusa hybrid 2.

Materials and Methods

Seeds were sown on nursery in three different dates viz. 20th July, 5th August and 20th August during 2001 and transplanted exactly after one month. A planting ratio of 4:2 (4 female: 2 male) and a spacing of 60 x 45 cm were maintained. Flowering started in the middle of February and ceased in the middle of March in 2003 in all the three dates of sowing. Observations on insect pollinators were recorded for four days (as four replication) in each date of sowing at 10.00, 12.00, and 14.00 and 16.00 hours on four plants each from both parental lines. Honeybees with pollen in their corbiculae were recorded as pollen gatherers and bees without pollen loads were recorded as nectar collectors. From each *Apis* species, 480 bees were observed for their movements between parental lines viz., male to male, male to female, female to female and female to male while foraging on the bloom. Nectar content was measured using graduated thin capillary tube (0.5 µl size) and nectar sugar content was measured by following phenol-sulphuric acid method (ROBERTS, 1979). Data on bee population was square root transformed and data on percentage proportion of honeybee movement between parental lines were angular transformed before being analysed.

Results

The field was predominantly visited by *Apis* species (85.23%). Other pollinators include dipteran flies; syrphid flies, moths and butterflies contributed 14.77% (Table I). Among *Apis* species, *Apis dorsata*, *Apis mellifera*, *Apis cerana indica* and *Apis florea* constituted 28.23, 26.32, 24.20 and 21.23% respectively. With respect to foraging modes, pollen gatherers (56.06%) found to be more predominant than nectar collectors (43.94%).

Table I

Pollinators of cauliflower hybrid	
Kind	Proportion (%)
A. Insects	
<i>Apis</i> species	85.23
Other insects	14.77
Honeybees	
<i>Apis dorsata</i>	28.23
<i>Apis mellifera</i>	26.32
<i>Apis cerana</i>	24.2
<i>Apis florea</i>	21.23
B. Foraging modes	
Pollen gatherers	56.06
Nectar collectors	43.94

Table II

Effect of different sowing dates and weather factors on abundance of *Apis* during peak flowering period

Sowing dates	Peak flowering period 2002	Duration (days)	Number of <i>Apis</i> per 64 plants*		Weather parameters			
					Temperature (°C)			RH%
			CC**	1-3-18-19***	Max.	Min.	Average	
July 20th	10/02 to 23/02	14	3306	3347	22.8	8.7	15.7	94.6
August 5th	20/02 to 03/03	12	3196	3207	23.1	10.6	16.9	91.5
August 20th	25/02 to 11/03	16	2817	2893	24.2	10.9	17.5	89.0

* Observed for 2 minutes per plant; ** Female parent; *** Male parent

Table III

Number of gatherers per plant during 2 minutes

Factors	Pollen gatherers	Nectar collectors
Sowing Dates		
July 20th	8.2 (2.91)	4.8 (2.28)
August 5th	8.11 (2.88)	4.44 (2.19)
August 20th	6.84 (2.68)	4.31 (2.17)
CD (P=0.05)	0.07	0.02
Hours of Day		
10.00	6.23 (2.55)	4.39 (2.19)
12.00	8.15 (2.90)	4.49 (2.21)
14.00	8.24 (3.01)	4.50 (2.21)
16.00	7.55 (2.82)	4.61 (2.24)
CD (P=0.05)	0.08	NS
Parental Lines		
CC	7.6 (2.80)	4.54 (2.22)
1-3-18-19	7.83 (2.84)	4.47 (2.21)
CD (P=0.05)	NS	NS
Honeybees		
<i>Apis dorsata</i>	10.16 (3.24)	5.56 (2.45)
<i>Apis mellifera</i>	8.68 (3.01)	4.85 (2.30)
<i>Apis cerana</i>	7.16 (2.74)	4.12 (2.13)
<i>Apis florea</i>	4.86 (2.30)	3.5 (1.98)
CD (P=0.05)	0.08	0.05

The number of pollen gatherers and nectar collecting bees per plant per two minutes are presented in table III. There was significant difference among different dates of sowing with regard to pollen gatherers. July 20th sown crop received more bee visits (8.20) than August 5th sown crop (8.11) and followed by August 20th sown crop (6.84). Pollen gatherers visited more at 14.00 (8.24) followed by 12.00 (8.15), 16.00 (7.55) and 10.00 hour (6.23). Pollen gatherers showed no significant difference between male and female parental lines. Among bees, *Apis dorsata* (10.16) visited more than *Apis mellifera* (8.68), *Apis cerana indica* (7.16) and *Apis florea* (4.86). Nectar collectors showed significant difference among different sowing dates and different bees. July 20th sown crop received more visits (4.80) than August 5th (4.44) and August 20th sown crop (4.31). Among bees, *Apis dorsata* (5.56) visited more than *Apis mellifera* (4.85), *Apis cerana indica* (4.12) and *Apis florea* (3.5). Between parental lines and among different hours of the day there were no significant differences with regard to nectar collectors.

Direction of honeybee movements viz. male to male, male to female, female to female and female to male parent were random as they were not significantly different among them (Table IV).

Table IV

Movements of honeybees

Directions	Proportion (%)
Male to male	25.3 (30.20)
Male to female	24.35 (29.57)
Female to female	21.43 (27.06)
Female to male	28.92 (27.06)
CD (P=0.05)	NS

Figures in parenthesis were angular transformed values

Table V

Nectar content (μ l) and nectar sugar content (μ g) per flower of parental lines of Pusa hybrid 2 at different hours of day

Floral rewards	Hours of day	CC	1-3-18-19	Mean
Nectar content	10.00	0.168	0.171	0.170
"	12.00	0.145	0.142	0.144
"	14.00	0.133	0.128	0.131
"	16.00	0.113	0.110	0.111
	Mean	0.140	0.138	
CD (P=0.05)	Parental lines (P): NS			
	Hours of day (D): 0.01			
	P X D: NS			
Nectar sugar content	10.00	0.453	0.353	0.403
"	12.00	0.698	0.523	0.610
"	14.00	0.835	0.682	0.758
"	16.00	1.146	0.850	0.998
	Mean	0.783	0.602	
CD (P=0.05)	Parental lines (P): 0.09			
	Hours of day (D): 0.123			
	P X D: NS			

Nectar content between parental lines showed no significant difference but nectar sugar content differed significantly (Table V). Nectar sugar content was more in female parent (0.782 μ g) than male parent (0.602 μ g).

Discussion

Honeybees were found as predominant pollinators (85.23%) in the seed production plot. Earlier, SHARMA et al. (1974) reported that honeybees were the predominant pollinators (42.1%) of cauliflower. SINHA and CHAKRABARTI (1980) reported that honeybees constituted 79, 82.4 and 83.3% respectively in 3 consecutive years. KAKKAR and SHARMA (1991) observed that honeybees constituted 38.7% on cauliflower bloom. These reports supported the present observation. Among honeybees, population of *Apis dorsata* was more than other *Apis* species irrespective of foraging modes. This might be due to presence of more colonies in nature. With respect to foraging modes, pollen gatherers outnumbered nectar collectors. During February and March heavy build up of colony (breeding) after overwintering necessitates more pollen grains to feed the larvae. This resulted in a greater number of pollen gatherers than nectar collectors.

July 20th sown crop received more bee visits than August 5th and August 20th sown crop. Change in major weather factors such as temperature and RH might be responsible for this variation (Table V). Bee foraging activity is highly influenced by prevailing weather factors (SZABO, 1980; SIHAG and ARBOL, 1986; ARBOL, 1987). Pollen gatherers visited more at 12 to 14 hours that coincided with maximum opening of flowers (anthesis). SINHA and CHAKRABARTI (1980) reported maximum honeybee visits at 12 to 14 hours on cauliflower. This supported present result that maximum bee visits recorded at 12 to 14 hours.

There was no significant difference between parental lines with regard to attractiveness of bees. Honeybee movements between parental lines were random and non selective. FAULKNER (1974 and 1976) and FREE and WILLIAMS (1983) reported that bees were able to discriminate between male and female parental lines of Brussels sprouts. A honeybee movement between parental lines was in the ratio of 30:1. The possible reason was differences in height and flower colour variation and also unknown factors. But parental lines of Pusa hybrid 2 were similar in height, flower colour and nectar content except nectar sugar content. High nectar sugar content had no effect on bee preference towards female parent. Correlation and path analysis of bee activity and different environmental factors revealed that nectar sugar concentration had no direct effect on bee activity (SIHAG and ARBOL, 1986; ARBOL, 1987, 1998).

Acknowledgment

I acknowledge the help rendered by Dr. R.N. YADAV and Dr. S.C. RANA during this investigation.

REFERENCES

- Abrol, D.P. (1987), Interrelation and path coefficient analysis of environmental factors influencing pollination activity of *Apis dorsata* F. on *Prunus persica* L., *Tropical Ecology* 28, 147-154
- Abrol, D.P. (1998), Environmental factors influencing flight activity in honeybees, *Apis cerana indica* and *Apis mellifera* (Hymenoptera: Apidae), *Indian Bee Journal* 60, 71-75
- Adlakha, R.L. and H.S.Dhaliwal (1979), Insect pollination of seed cauliflower (*Brassica oleracea* var. *botrytis*) with particular reference to the role of honeybees, *Indian Bee Journal*, 41, 13-16
- Faulkner, G.J. (1974), Factors affecting field scale production of seed of F1 hybrid Brussels sprout, *Annals of Applied Biology* 77, 181-190
- Faulkner, G.J. (1976), Honeybee behaviour as affected by plant height and flower colour variation in Brussels sprouts, *Journal of Apicultural Research* 15, 15-18
- Free, J.B. and I.H.Williams (1983), Foraging behaviour of honeybees and bumble bees on Brussels sprouts grown to produce hybrid seed, *Journal of Apicultural Research* 22, 94-97
- Kakar, K.L. (1981), Foraging behaviour of insect pollination of cauliflower bloom, *Indian Journal of Ecology* 8, 126-130
- Kakkar, K.L. and P.L. Sharma (1991), Studies on the role of honeybee, *Apis cerana indica* F. in the pollination of cauliflower, *Brassica oleracea* var. *botrytis*. *Indian Journal of Entomology*, 53, 66-69
- Raula, T.S. (1972), Pollination studies in cauliflower (*Brassica oleracea* var. *botrytis*), *Journal of Research Punjab Agricultural University* 9, 580-585
- Sharma, A.K., H.S.Dhaliwal and K.L.Kakar (1974), Insect visitors and pollinators of cauliflower (*Brassica oleracea* var. *botrytis*) seed crop bloom, *Himachal Journal of Agricultural Research* 2, 74-78
- Sihag, R.C. and D.P.Abrol (1986), Correlation and path analysis of environmental factors influencing flight activity of *Apis florea* F., *Journal of Apicultural Research* 25, 202-208
- Sinha, S.N. and A.K. Chakrabarti (1980), Bee pollination and its impact on cauliflower seed production, Proceedings of the Second International Conference on Apiculture in Tropical Climates, NewDelhi, February 29th - march 4th, 1980, 1983, 513-527
- Szabo, T.I. (1980), Effects of weather factors on honeybee activity and colony weight gain, *Journal of Apicultural Research* 19, 164-171